

5

ANTIANGIOGENIC AGENTS

10

Im917

FIELD OF THE INVENTION

15

The present invention relates to treating disease states characterized by abnormal cell mitosis and or abnormal angiogenesis. More particularly, the present invention relates to certain analogs of 2-methoxyestradiol (2ME2) and their effect on diseases characterized by abnormal cell mitosis and/or abnormal angiogenesis.

BACKGROUND OF THE INVENTION

20

As used herein, the term "angiogenesis" means the generation of new blood vessels into a tissue or organ. Under normal physiological conditions, humans or animals only undergo angiogenesis in very specific restricted situations. For example, angiogenesis is normally observed in wound healing, fetal and embryonal development and formation of the corpus luteum, endometrium and placenta. The control of angiogenesis is a highly regulated system of angiogenic stimulators and inhibitors. The control of angiogenesis has been found to be altered in certain disease states and, in many cases, the pathological damage associated with the disease is related to the uncontrolled angiogenesis.

25

30

35

Both controlled and uncontrolled angiogenesis are thought to proceed in a similar manner. Endothelial cells and pericytes, surrounded by a basement membrane, form capillary blood vessels. Angiogenesis begins with the erosion of the basement membrane by enzymes released by endothelial cells and leukocytes. The endothelial cells, which line the lumen of blood vessels, then protrude through the basement membrane. Angiogenic stimulants induce the endothelial cells to migrate through the eroded basement

membrane. The migrating cells form a "sprout" off the parent blood vessel, where the endothelial cells undergo mitosis and proliferate. The endothelial sprouts merge with each other to form capillary loops, creating the new blood vessel. In the disease state, prevention of angiogenesis could avert the damage caused by the invasion of the new microvascular system.

Persistent, unregulated angiogenesis occurs in a multiplicity of disease states, tumor metastasis and abnormal growth by endothelial cells and supports the pathological damage seen in these conditions. The diverse pathological states created due to unregulated angiogenesis have been grouped together as angiogenic dependent or angiogenic associated diseases. Therapies directed at control of the angiogenic processes could lead to the abrogation or mitigation of these diseases.

One example of a disease mediated by angiogenesis is ocular neovascular disease. This disease is characterized by invasion of new blood vessels into the structures of the eye such as the retina or cornea. It is the most common cause of blindness and is involved in approximately twenty eye diseases. In age-related macular degeneration, the associated visual problems are caused by an ingrowth of chorioidal capillaries through defects in Bruch's membrane with proliferation of fibrovascular tissue beneath the retinal pigment epithelium. Angiogenic damage is also associated with diabetic retinopathy, retinopathy of prematurity, corneal graft rejection, neovascular glaucoma and retrolental fibroplasia. Other diseases associated with corneal neovascularization include, but are not limited to, epidemic keratoconjunctivitis, Vitamin A deficiency, contact lens overwear, atopic keratitis, superior limbic keratitis, pterygium keratitis sicca, sjogrens, acne rosacea, phlyctenulosis, syphilis, *Mycobacteria* infections, lipid degeneration, chemical burns, bacterial ulcers, fungal ulcers, *Herpes simplex* infections, *Herpes zoster* infections, protozoan infections, Kaposi sarcoma, Mooren ulcer, Terrien's marginal degeneration, marginal keratolysis, rheumatoid arthritis, systemic lupus, polyarteritis, trauma, Wegeners sarcoidosis, Scleritis, Steven's Johnson disease, periphigoid radial keratotomy, and corneal graft rejection.

Diseases associated with retinal/choroidal neovascularization include, but are not limited to, diabetic

retinopathy, macular degeneration, sickle cell anemia, sarcoid, syphilis, pseudoxanthoma elasticum, Pagets disease, vein occlusion, artery occlusion, carotid obstructive disease, chronic uveitis/vitritis, mycobacterial infections, Lyme's disease, systemic lupus erythematosus, retinopathy of prematurity, Eales disease, Bechets disease, infections causing a retinitis or choroiditis, presumed ocular histoplasmosis, Bests disease, myopia, optic pits, Stargarts disease, pars planitis, chronic retinal detachment, hyperviscosity syndromes, toxoplasmosis, trauma and post-laser complications. Other diseases include, but are not limited to, diseases associated with rubeosis (neovasculariation of the angle) and diseases caused by the abnormal proliferation of fibrovascular or fibrous tissue including all forms of proliferative vitreoretinopathy.

Another disease in which angiogenesis is believed to be involved is rheumatoid arthritis. The blood vessels in the synovial lining of the joints undergo angiogenesis. In addition to forming new vascular networks, the endothelial cells release factors and reactive oxygen species that lead to pannus growth and cartilage destruction. The factors involved in angiogenesis may actively contribute to, and help maintain, the chronically inflamed state of rheumatoid arthritis.

Factors associated with angiogenesis may also have a role in osteoarthritis. The activation of the chondrocytes by angiogenic-related factors contributes to the destruction of the joint. At a later stage, the angiogenic factors would promote new bone formation. Therapeutic intervention that prevents the bone destruction could halt the progress of the disease and provide relief for persons suffering with arthritis.

Chronic inflammation may also involve pathological angiogenesis. Such disease states as ulcerative colitis and Crohn's disease show histological changes with the ingrowth of new blood vessels into the inflamed tissues. Bartonellosis, a bacterial infection found in South America, can result in a chronic stage that is characterized by proliferation of vascular endothelial cells. Another pathological role associated with angiogenesis is found in atherosclerosis. The plaques formed within the lumen of blood vessels have been shown to have angiogenic stimulatory activity.

One of the most frequent angiogenic diseases of childhood is the hemangioma. In most cases, the tumors are benign

and regress without intervention. In more severe cases, the tumors progress to large cavernous and infiltrative forms and create clinical complications. Systemic forms of hemangiomas, the hemangiomatoses, have a high mortality rate. Therapy-resistant hemangiomas exist that cannot be treated with therapeutics currently in use.

Angiogenesis is also responsible for damage found in hereditary diseases such as Osler-Weber-Rendu disease, or hereditary hemorrhagic telangiectasia. This is an inherited disease characterized by multiple small angiomas, tumors of blood or lymph vessels. The angiomas are found in the skin and mucous membranes, often accompanied by epistaxis (nosebleeds) or gastrointestinal bleeding and sometimes with pulmonary or hepatic arteriovenous fistula.

Angiogenesis is prominent in solid tumor formation and metastasis. Angiogenic factors have been found associated with several solid tumors such as rhabdomyosarcomas, retinoblastoma, Ewing sarcoma, neuroblastoma, and osteosarcoma. A tumor cannot expand without a blood supply to provide nutrients and remove cellular wastes. Tumors in which angiogenesis is important include solid tumors, and benign tumors such as acoustic neuroma, neurofibroma, trachoma and pyogenic granulomas. Prevention of angiogenesis could halt the growth of these tumors and the resultant damage to the animal due to the presence of the tumor.

It should be noted that angiogenesis has been associated with blood-born tumors such as leukemias, any of various acute or chronic neoplastic diseases of the bone marrow in which unrestrained proliferation of white blood cells occurs, usually accompanied by anemia, impaired blood clotting, and enlargement of the lymph nodes, liver, and spleen. It is believed that angiogenesis plays a role in the abnormalities in the bone marrow that give rise to leukemia-like tumors.

Angiogenesis is important in two stages of tumor metastasis. The first stage where angiogenesis stimulation is important is in the vascularization of the tumor which allows tumor cells to enter the blood stream and to circulate throughout the body. After the tumor cells have left the primary site, and have settled into the secondary, metastasis site, angiogenesis must occur before the new tumor can grow and expand. Therefore, prevention of

angiogenesis could lead to the prevention of metastasis of tumors and possibly contain the neoplastic growth at the primary site.

Knowledge of the role of angiogenesis in the maintenance and metastasis of tumors has led to a prognostic indicator for breast cancer. The amount of neovascularization found in the primary tumor was determined by counting the microvessel density in the area of the most intense neovascularization in invasive breast carcinoma. A high level of microvessel density was found to correlate with tumor recurrence. Control of angiogenesis by therapeutic means could possibly lead to cessation of the recurrence of the tumors.

Angiogenesis is also involved in normal physiological processes such as reproduction and wound healing. Angiogenesis is an important step in ovulation and also in implantation of the blastula after fertilization. Prevention of angiogenesis could be used to induce amenorrhea, to block ovulation or to prevent implantation by the blastula.

In wound healing, excessive repair or fibroplasia can be a detrimental side effect of surgical procedures and may be caused or exacerbated by angiogenesis. Adhesions are a frequent complication of surgery and lead to problems such as small bowel obstruction.

Several kinds of compounds have been used to prevent angiogenesis. Taylor et al. have used protamine to inhibit angiogenesis, see Taylor et al., *Nature* 297:307 (1982). The toxicity of protamine limits its practical use as a therapeutic. Folkman et al. have disclosed the use of heparin and steroids to control angiogenesis. See Folkman et al., *Science* 221:719 (1983) and U.S. Patent Nos. 5,001,116 and 4,994,443. Steroids, such as tetrahydrocortisol, which lack gluco and mineral corticoid activity, have been found to be angiogenic inhibitors.

Other factors found endogenously in animals, such as a 4 kDa glycoprotein from bovine vitreous humor and a cartilage derived factor, have been used to inhibit angiogenesis. Cellular factors such as interferon inhibit angiogenesis. For example, interferon α or human interferon β has been shown to inhibit tumor-induced angiogenesis in mouse dermis stimulated by human neoplastic cells. Interferon β is also a potent inhibitor of angiogenesis induced by allogeneic spleen cells. See Sidky et al.,

Cancer Research 47:5155-5161 (1987). Human recombinant α interferon (alpha/A) was reported to be successfully used in the treatment of pulmonary hemangiomatosis, an angiogenesis-induced disease. See White et al., *New England J. Med.* 320:1197-1200 (1989).

Other agents which have been used to inhibit angiogenesis include ascorbic acid ethers and related compounds. See Japanese Kokai Tokkyo Koho No. 58-131978. Sulfated polysaccharide DS 4152 also shows angiogenic inhibition. See Japanese Kokai Tokkyo Koho No. 63-119500. A fungal product, fumagillin, is a potent angiostatic agent *in vitro*. The compound is toxic *in vivo*, but a synthetic derivative, AGM 12470, has been used *in vivo* to treat collagen II arthritis. Fumagillin and O-substituted fumagillin derivatives are disclosed in EPO Publication Nos. 0325199A2 and 0357061A1. Folkman et al., described several proteins derived from endogenous proteins including angiostatin and endostatin. (See, for example, U.S. Patent No. 6,024,688 and U.S. Patent No. 5,854,205 which are incorporated in their entirety) D'Amato et al., described 2-methoxyestradiol and derivatives of 2-methoxyestradiol in U.S. Patent Nos. 5,504,074 and 5,661,143 which are incorporated herein by reference entirety.

The above compounds are either topical or injectable therapeutics. Therefore, there are drawbacks to their use as a general angiogenic inhibitor and lack adequate potency. For example, in prevention of excessive wound healing, surgery on internal body organs involves incisions in various structures contained within the body cavities. These wounds are not accessible to local applications of angiogenic inhibitors. Local delivery systems also involve frequent dressings which are impracticable for internal wounds, and increase the risk of infection or damage to delicate granulation tissue for surface wounds.

Thus, a method and composition are needed that are capable of inhibiting angiogenesis and which are easily administered. A simple and efficacious method of treatment would be through the oral route. If an angiogenic inhibitor could be given by an oral route, the many kinds of diseases discussed above, and other angiogenic dependent pathologies, could be treated easily. The optimal dosage could be distributed in a form that the patient could self-administer.

Other diseases are also characterized by an abnormal balance between cellular mitosis and apoptosis. One of these diseases is osteoporosis. Osteoporosis is characterized by a reduction in the bone mass of the skeleton which leads to skeletal fragility and an increased risk of fracture. In humans, the most common sites of fracture are found in the forearm, the vertebrae and the hip bones. Osteoporosis and its attendant fractures are a major cause of morbidity and mortality and lead to increased health costs for care.

In treating osteoporosis the main objective is to prevent fractures by stopping the loss of skeletal integrity. A variety of different therapies have been tried to achieve this objective, such as calcium, Vitamin D supplements and hormone replacement. Calcitonin has been used to improve bone mineral density at all bone sites. Bisphosphonates are an important group of therapeutic agents used for treatment of osteoporosis. They act by inhibiting bone resorption and increase bone density. Cyclical etidronate treatment aids in decreasing vertebral fractures, as does hormone replacement therapy and calcitonin. Alendronate has been shown to decrease the risk of symptomatic fractures of the forearm, spine and hip.

None of these treatments have proven to be effective in large numbers of osteoporotic patients. Additionally, the currently used therapies have unwanted side effects that create compliance and tolerance problems in treatment regimens. The most common adverse events with cyclical etidronate and alendronate are gastrointestinal disturbances. Esophagitis has also been a complication of therapies with alendronate. Cyclical etidronate has been shown to lead to focal osteomalacia. Hormone replacement therapies lead to estrogen effects such as uterine hypertrophy, and a potential for stimulation of estrogen-sensitive tumors leading to complications such as breast cancer.

What is needed are safe and effective treatments that do not create unwanted side effects.

2-Methoxyestradiol is an endogenous metabolite of estradiol (E2) that has potent anti-proliferative activity and induces apoptosis in a wide variety of tumor and non-tumor cell lines. When administered orally, it exhibits anti-tumor and anti-proliferative activity with little or no toxicity. *In vitro* data suggests that 2-methoxyestradiol does not engage the estrogen receptor for its anti-

proliferative activity and is not estrogenic over a wide range of concentrations, as assessed by estrogen dependant MCF-7 cell proliferation. However, the presence of demethylases *in vivo* may metabolize this compound to 2-hydroxyestradiol, which has been shown to be estrogenic by several approaches. What is needed is a means to improve the bioavailability of estradiol or 2-methoxyestradiol and to reduce the formation of estrogenic 2-methoxyestradiol metabolites. What is also needed is a means to modify estradiol or 2-methoxyestradiol in such a way that the molecule can not be converted into an uterotrophic derivative.

SUMMARY OF THE INVENTION

The present invention provides certain analogs of 2-methoxyestradiol that are effective in treating diseases characterized by abnormal mitosis and/or abnormal angiogenesis. Specifically the present invention relates to analogs of 2-methoxyestradiol that have been modified at the 2 position and the 16 position. Compounds within the general formulae that inhibit cell proliferation are preferred. Preferred compositions may also exhibit a change (increase or decrease) in estrogen receptor binding, improved absorption, transport (e.g. through blood-brain barrier and cellular membranes), biological stability, or decreased toxicity. The invention also provides compounds useful in the method, as described by the general formulae of the claims.

A mammalian disease characterized by undesirable cell mitosis, as defined herein, includes but is not limited to excessive or abnormal stimulation of endothelial cells (e.g., atherosclerosis), solid tumors and tumor metastasis, benign tumors, for example, hemangiomas, acoustic neuromas, neurofibromas, trachomas, and pyogenic granulomas, vascular malfunctions, abnormal wound healing, inflammatory and immune disorders, Bechet's disease, gout or gouty arthritis, abnormal angiogenesis accompanying: rheumatoid arthritis, psoriasis, diabetic retinopathy, and other ocular angiogenic diseases such as retinopathy of prematurity (retrolental fibroplastic), macular degeneration, corneal graft rejection, neovascular glaucoma and Osler Weber syndrome. Other undesired angiogenesis involves normal processes including ovulation and implantation of a blastula. Accordingly, the compositions described above can be used to block

ovulation and implantation of a blastula or to block menstruation (induce amenorrhea).

Since 2-methoxyestradiol is metabolized to a much less active metabolite, the present invention adds steric bulk and/or modification of electrostatic characteristics at position 16 of 2-methoxyestradiol for retarding or preventing interaction of 17 β -hydroxysteroid dehydrogenases and co-factor NADP⁺ on this substrate. Addition of steric bulk and/or modification of electrostatic characteristics at position 16 of 2-methoxyestradiol may retard or prevent glucuronidation. It is believed that retardation or prevention of these two metabolic deactivation pathways prolongs the serum lifetime of 2-methoxyestradiol and other estrogenic compounds while retaining the desired anti-angiogenic and anti-tumor activity.

Aside from preventing the possible metabolism of 2ME2 to 2ME1, which may occur by making these steroids poor substrates for 17B-HSD (by either steric and / or electronic effects), it is not possible for these analogs to undergo the demethylation known to occur with 2ME2 since there is no methyl ether group at that position. This is desirable since it has been demonstrated that 2-hydroxyestradiol (the product of demethylation of 2ME2) has estrogenic activity.

Also disclosed is a method for modifying the methyl ether of 2-methoxyestradiol so that it can not be a substrate for demethylase and the resulting compounds.

Other features and advantages of the invention will be apparent from the following description of preferred embodiments thereof.

BRIEF DESCRIPTION OF THE DRAWINGS

Fig. 1 depicts: I. colchicine, 2-methoxyestradiol and combretastatin A-4, and II. various estradiol derivatives comprising colchicine (a-c) or combretastatin A-4 (d) structural motifs as described below.

DETAILED DESCRIPTION OF THE INVENTION

As described below, compounds that are useful in accordance with the invention include novel estradiol derivatives that exhibit anti-mitotic, anti-angiogenic and anti-tumor properties.

Specific compounds according to the invention are described below. Preferred compounds of the invention are estradiol derivatives modified at either the 2 or 16 positions. Those skilled in the art will appreciate that the invention extends to other compounds within the formulae given in the claims below, having the described characteristics. These characteristics can be determined for each test compound using the assays detailed below and elsewhere in the literature.

Without wishing to be bound to specific mechanisms or theory, it appears that certain compounds that are known to exhibit anti-mitotic properties such as colchicine and combretastatin A-4 share certain structural similarities with estradiol. Fig. 1 illustrates the molecular formulae of estradiol, colchicine, combretastatin A-4, and improved estradiol derivatives that exhibit anti-mitotic, anti-angiogenic and anti-tumor properties. Molecular formulae are drawn and oriented to emphasize structural similarities between the ring structures of colchicine, combretastatin A-4, estradiol, and certain estradiol derivatives. Estradiol derivatives are made by incorporating colchicine or combretastatin A-4 structural motifs into the steroidal backbone of estradiol.

Figure 1, part I, depicts the chemical formulae of colchicine, 2-methoxyestradiol and combretastatin A-4. Figure 1, part II a-d, illustrates estradiol derivatives that comprise structural motifs found in colchicine or combretastatin A-4. For example, part II a-c shows estradiol derivatives with an A and/or B ring expanded from six to seven carbons as found in colchicine and part IId depicts an estradiol derivative with a partial B ring as found in combretastatin A-4. Each C ring of an estradiol derivative, including those shown in Figure 1, may be fully saturated as found in 2-methoxyestradiol. R₁₋₆ represent a subset of the substitution groups found in the claims. Each R₁-R₆ can independently be defined as -R₁, OR₁, -OCOR₁₁-SR₁, -F, -NHR₂, -Br, -I, or -C≡CH.

2-Methoxyestradiol is an endogenous metabolite of estradiol that has potent anti-proliferative activity and induces apoptosis in a wide variety of tumor and non-tumor cell lines. When administered orally, it exhibits anti-tumor and anti-proliferative activity with little or no toxicity. 2-Methoxyestradiol is metabolized

5 to a much less active metabolite, 2-methoxyestrone as indicated by
in vitro and in vivo results. Although not wishing to be bound by
theory, it is believed that this metabolite is formed through the same
enzymatic pathway as estrone is formed from estradiol. Although
10 not wishing to be bound by theory, it is believed that the enzymes
responsible for this reversible reaction on estradiol are the 17 β -
hydroxysteroid dehydrogenases (17 β -HSD) and NADP+ co-factor
(Han *et al.*, *J. Biol. Chem.* 275:2, 1105-1111 (Jan. 12, 2000) and
other references cited earlier). Each of the four members of this
15 enzyme family, types 1, 2, 3, and 4, have distinct activity. It appears
that 17 β -HSD type 1 catalyzes the reductive reaction (estrone to
estradiol), while 17 β -HSD type 2 catalyzes the oxidation reaction
(estradiol to estrone), and type 3 catalyzes 4-androstenedione to
testosterone. An additional metabolic deactivation pathway results in
glucuronidation of 2-methoxyestradiol.

20 Since 2-methoxyestradiol is metabolized to a much less
active metabolite, the present invention adds steric bulk and/or
modification of electrostatic characteristics at position 16 of 2-
methoxyestradiol for retarding or preventing interaction of the
family of 17 β -hydroxysteroid dehydrogenases and co-factor NADP+
on this substrate. Addition of steric bulk and/or modification of
electrostatic characteristics at position 16 of 2-methoxyestradiol also
retards or prevents glucuronidation. It is believed that retardation or
prevention of these two metabolic deactivation pathways prolongs
25 the serum lifetime of 2-methoxyestradiol and other estradiol
derivatives while retaining the desired anti-angiogenic and anti-tumor
activity.

30 Aside from preventing the possible metabolism of
2ME2 to 2ME1, which may occur by making these steroids poor
substrates for 17 β -HSD (by either steric and / or electronic effects),
it is not possible for these analogs to undergo the demethylation
known to occur with 2ME2 since there is no methyl ether group at
that position. This is desirable since it has been demonstrated that 2-
hydroxyestradiol (the product of demethylation of 2ME2) has
35 estrogenic activity.

In another embodiment of the invention, estradiol
derivatives are modified at the 2 position.

Anti-Proliferative Activity *In Situ*

Anti-proliferative activity is evaluated *in situ* by testing the ability of an improved estradiol derivative to inhibit the proliferation of new blood vessel cells (angiogenesis). A suitable assay is the chick embryo chorioallantoic membrane (CAM) assay described by Crum et al. *Science* 230:1375 (1985). See also, U.S. Patent 5,001,116, hereby incorporated by reference, which describes the CAM assay. Briefly, fertilized chick embryos are removed from their shell on day 3 or 4, and a methylcellulose disc containing the drug is implanted on the chorioallantoic membrane. The embryos are examined 48 hours later and, if a clear avascular zone appears around the methylcellulose disc, the diameter of that zone is measured. Using this assay, a 100 mg disk of the estradiol derivative 2-methoxyestradiol was found to inhibit cell mitosis and the growth of new blood vessels after 48 hours. This result indicates that the anti-mitotic action of 2-methoxyestradiol can inhibit cell mitosis and angiogenesis.

Anti-Proliferative Activity *In Vitro*

The process by which 2ME₂ affects cell growth remains unclear, however, a number of studies have implicated various mechanisms of action and cellular targets. 2ME₂ induced changes in the levels and activities of various proteins involved in the progression of the cell cycle. These include cofactors of DNA replication and repair, e.g., proliferating cell nuclear antigen (PCNA) (Klauber, N., Parangi, S., Flynn, E., Hamel, E. and D'Amato, R.J. (1997), Inhibition of angiogenesis and breast cancer in mice by the microtubule inhibitors 2-methoxyestradiol and Taxol., *Cancer Research* 57, 81-86; Lottering, M-L., de Kock, M., Viljoen, T.C., Grobler, C.J.S. and Seegers, J.C. (1996) 17 β -estradiol metabolites affect some regulators of the MCF-7 cell cycle. *Cancer Letters* 110, 181-186); cell division cycle kinases and regulators, e.g., p34^{cdc2} and cyclin B (Lottering *et al.* (1996); Attalla, H., Mäkelä, T.P., Adlercreutz, H. and Andersson, L.C. (1996) 2-methoxyestradiol arrests cells in mitosis without depolymerizing tubulin. *Biochemical and Biophysical Research Communications* 228, 467-473; Zoubine, M.N., Weston, A.P., Johnson, D.C., Campbell, D.R. and Banerjee, S.K. (1999) 2-Methoxyestradiol-induced growth suppression and

lethality in estrogen- responsive MCF-7 cells may be mediated by
 down regulation of p34cdc2 and cyclin B1 expression. *Int J Oncol*
 15, 639-646); transcription factor modulators, e.g., SAPK/JNK (Yue,
 T-L., Wang, X., Louden, C.S., Gupta, L.S., Pillarisetti, K., Gu, J-L.,
 Hart, T.K., Lysko, P.G. and Feuerstein, G.Z. (1997) 2-
 methoxyestradiol, an endogenous estrogen metabolite induces
 apoptosis in endothelial cells and inhibits angiogenesis: Possible role
 for stress-activated protein kinase signaling pathway and fas
 expression. *Molecular Pharmacology* 51, 951-962; Attalla, H.,
 Westberg, J.A., Andersson, L.C., Aldercreutz, H. and Makela, T.P.
 (1998) 2-Methoxyestradiol-induced phosphorylation of bcl-2:
 uncoupling from JNK/SAPK activation. *Biochem and Biophys Res*
Commun 247, 616-619); and regulators of cell arrest and apoptosis,
 e.g., tubulin (D'Amato, R.J., Lin, C.M., Flynn, E., Folkman, J. and
 Hamel, E. (1994) 2-Methoxyestradiol, and endogenous mammalian
 metabolite, inhibits tubulin polymerization by interacting at the
 colchicine site. *Proc. Natl. Acad. Sci. USA* 91, 3964-3968; Hamel,
 E., Lin, C.M., Flynn, E. and D'Amato, R.J. (1996) Interactions of 2-
 methoxyestradiol, and endogenous mammalian metabolite, with
 unpolymerized tubulin and with tubulin polymers. *Biochemistry* 35,
 1304-1310), p21^{WAF1/CIP1} (Mukhopadhyay, T. and Roth, J.A. (1997)
 Induction of apoptosis in human lung cancer cells after wild-type p53
 activation by methoxyestradiol. *Oncogene* 14, 379-384), bcl-2 and
 FAS (Yue *et al.* (1997); Attalla *et al.* (1998)), and p53 (Kataoka, M.,
 Schumacher, G., Cristiano, R.J., Atkinson, E.N., Roth, J.A. and
 Mukhopadhyay, T. (1998) An agent that increases tumor suppressor
 transgene product coupled with systemic transgene delivery inhibits
 growth of metastatic lung cancer *in vivo*. *Cancer Res* 58, 4761-4765;
 Mukhopadhyay *et al.* (1997); Seegers, J.C., Lottering, M-L., Grobler
 C.J.S., van Papendorp, D.H., Habbersett, R.C., Shou, Y. and Lehnert
 B.E. (1997) The mammalian metabolite, 2-methoxyestradiol, affects
 p53 levels and apoptosis induction in transformed cells but not in
 normal cells. *J. Steroid Biochem. Molec.Biol.* 62, 253-267). The
 effects on the level of cAMP, calmodulin activity and protein
 phosphorylation may also be related to each other. More recently,
 2ME2 was shown to upregulate Death Receptor 5 and caspase 8 in
 human endothelial and tumor cell lines (LaVallee, T.M., Hembrough,
 W.A., Williams, M.S., Zhan, X.H., Pribluda, V.S., Papathanassiou, A.,

and Green, S.J. 2-Methoxyestradiol upregulates DR5 and induces apoptosis independently of p53. (Submitted for publication)). All cellular targets described above are not necessarily mutually exclusive to the inhibitory effects of 2ME₂ in actively dividing cells.

5 The high affinity binding to SHBG has been mechanistically associated to its efficacy in a canine model of prostate cancer, in which signaling by estradiol and 5 α -androstan-3 α ,17 β -diol were inhibited by 2ME₂ (Ding, V.D., Moller, D.E., Feeney, W.P., Didolkar, V., Nakhla, A.M., Rhodes, L., Rosner, W. and Smith, R.G. 10 (1998) Sex hormone-binding globulin mediates prostate androgen receptor action via a novel signaling pathway. *Endocrinology* 139, 213-218).

15 The more relevant mechanism described above have been extensively discussed in Victor S. Pribluda, Theresa M. LaVallee and Shawn J. Green, 2-methoxyestradiol: *a novel endogenous chemotherapeutic and antiangiogenic* in The New Angiotherapy, Tai-Ping Fan and Robert Auerbach eds., Human Press Publisher.

20 Assays relevant to the mechanisms of action and activity are well-known in the art. For example, anti-mitotic activity mediated by effects on tubulin polymerization activity can be evaluated by testing the ability of an estradiol derivative to inhibit tubulin polymerization and microtubule assembly *in vitro*. Microtubule assembly is followed in a Gilford recording spectrophotometer (model 250 or 2400S) equipped with electronic temperature controllers. A reaction mixture (all concentrations refer to a final reaction volume of 0.25 μ l) contains 1.0M monosodium glutamate (pH 6.6), 1.0mg/ml (10 μ M) tubulin, 1.0 mM MgCl₂, 4% (v/v) dimethylsulfoxide and 20-75 μ M of a composition to be tested. 25 The 0.24ml reaction mixtures are incubated for 15 min. at 37 $^{\circ}$ C and then chilled on ice. After addition of 10 μ l 2.5mM GTP, the reaction mixture is transferred to a cuvette at 0 $^{\circ}$ C, and a baseline established. 30 At time zero, the temperature controller of the spectrophotometer is set at 37 $^{\circ}$ C. Microtubule assembly is evaluated by increased turbidity at 350 nm. Alternatively, inhibition of microtubule assembly can be followed by transmission electron microscopy as described in 35 Example 2 below.

Other such assays include counting of cells in tissue culture plates or assessment of cell number through metabolic assays or incorporation into DNA of labeled (^3H -thymidine) or immuno-reactive (BrdU) nucleotides. In addition, antiangiogenic activity may be evaluated through endothelial cell migration, endothelial cell tubule formation, or vessel outgrowth in ex-vivo models such as rat aortic rings.

Indications

The invention can be used to treat any disease characterized by abnormal cell mitosis. Such diseases include, but are not limited to: abnormal stimulation of endothelial cells (e.g., atherosclerosis), solid tumors and tumor metastasis, benign tumors, for example, hemangiomas, acoustic neuromas, neurofibromas, trachomas, and pyogenic granulomas, vascular malfunctions, abnormal wound healing, inflammatory and immune disorders, Bechet's disease, gout or gouty arthritis, abnormal angiogenesis accompanying: rheumatoid arthritis, psoriasis, diabetic retinopathy, and other ocular angiogenic diseases such as retinopathy of prematurity (retrolental fibroplastic), macular degeneration, corneal graft rejection, neurovascular glaucoma and Oster Webber syndrome.

In addition, the invention can be used to treat a variety of post-menopausal symptoms, including osteoporosis, cardiovascular disease, Alzheimer's disease, to reduce the incidence of strokes, and as an alternative to prior estrogen replacement therapies. The compounds of the present invention can work by estrogenic and non-estrogenic biochemical pathways.

Improved Estradiol Derivative Synthesis

Known compounds that are used in accordance with the invention and precursors to novel compounds according to the invention can be purchased, e.g., from Sigma Chemical Co., St. Louis, Steraloids and Research Plus. Other compounds according to the invention can be synthesized according to known methods from publicly available precursors.

The chemical synthesis of estradiol has been described (Eder, V. et al., *Ber* 109, 2948 (1976); Oppolzer, D.A. and Roberts,

5 DA. *Helv. Chim. Acta.* 63, 1703, (1980)). Synthetic methods for making seven-membered rings in multi-cyclic compounds are known (Nakamura, T. et al. *Chem. Pharm. Bull.* 10, 281 (1962); Sunagawa, G. et al. *Chem. Pharm. Bull.* 9, 81 (1961); Van Tamelen, E. E. et al. *Tetrahedron* 14, 8-34 (1961); Evans, D. E. et al. *JACS* 103, 5813 (1981)). Those skilled in the art will appreciate that the chemical synthesis of estradiol can be modified to include 7-membered rings by making appropriate changes to the starting materials, so that ring closure yields seven-membered rings. Estradiol or estradiol derivatives can be modified to include appropriate chemical side groups according to the invention by known chemical methods (*The Merck Index*, 11th Ed., Merck & Co., Inc., Rahway, NJ USA (1989), pp. 583-584).

15
20
25
30
35
40
45
50
55
60
65
70
75
80
85
90
95
100
105
110
115
120
125
130
135
140
145
150
155
160
165
170
175
180
185
190
195
200
205
210
215
220
225
230
235
240
245
250
255
260
265
270
275
280
285
290
295
300
305
310
315
320
325
330
335
340
345
350
355
360
365
370
375
380
385
390
395
400
405
410
415
420
425
430
435
440
445
450
455
460
465
470
475
480
485
490
495
500
505
510
515
520
525
530
535
540
545
550
555
560
565
570
575
580
585
590
595
600
605
610
615
620
625
630
635
640
645
650
655
660
665
670
675
680
685
690
695
700
705
710
715
720
725
730
735
740
745
750
755
760
765
770
775
780
785
790
795
800
805
810
815
820
825
830
835
840
845
850
855
860
865
870
875
880
885
890
895
900
905
910
915
920
925
930
935
940
945
950
955
960
965
970
975
980
985
990
995
1000
1005
1010
1015
1020
1025
1030
1035
1040
1045
1050
1055
1060
1065
1070
1075
1080
1085
1090
1095
1100
1105
1110
1115
1120
1125
1130
1135
1140
1145
1150
1155
1160
1165
1170
1175
1180
1185
1190
1195
1200
1205
1210
1215
1220
1225
1230
1235
1240
1245
1250
1255
1260
1265
1270
1275
1280
1285
1290
1295
1300
1305
1310
1315
1320
1325
1330
1335
1340
1345
1350
1355
1360
1365
1370
1375
1380
1385
1390
1395
1400
1405
1410
1415
1420
1425
1430
1435
1440
1445
1450
1455
1460
1465
1470
1475
1480
1485
1490
1495
1500
1505
1510
1515
1520
1525
1530
1535
1540
1545
1550
1555
1560
1565
1570
1575
1580
1585
1590
1595
1600
1605
1610
1615
1620
1625
1630
1635
1640
1645
1650
1655
1660
1665
1670
1675
1680
1685
1690
1695
1700
1705
1710
1715
1720
1725
1730
1735
1740
1745
1750
1755
1760
1765
1770
1775
1780
1785
1790
1795
1800
1805
1810
1815
1820
1825
1830
1835
1840
1845
1850
1855
1860
1865
1870
1875
1880
1885
1890
1895
1900
1905
1910
1915
1920
1925
1930
1935
1940
1945
1950
1955
1960
1965
1970
1975
1980
1985
1990
1995
2000
2005
2010
2015
2020
2025
2030
2035
2040
2045
2050
2055
2060
2065
2070
2075
2080
2085
2090
2095
2100
2105
2110
2115
2120
2125
2130
2135
2140
2145
2150
2155
2160
2165
2170
2175
2180
2185
2190
2195
2200
2205
2210
2215
2220
2225
2230
2235
2240
2245
2250
2255
2260
2265
2270
2275
2280
2285
2290
2295
2300
2305
2310
2315
2320
2325
2330
2335
2340
2345
2350
2355
2360
2365
2370
2375
2380
2385
2390
2395
2400
2405
2410
2415
2420
2425
2430
2435
2440
2445
2450
2455
2460
2465
2470
2475
2480
2485
2490
2495
2500
2505
2510
2515
2520
2525
2530
2535
2540
2545
2550
2555
2560
2565
2570
2575
2580
2585
2590
2595
2600
2605
2610
2615
2620
2625
2630
2635
2640
2645
2650
2655
2660
2665
2670
2675
2680
2685
2690
2695
2700
2705
2710
2715
2720
2725
2730
2735
2740
2745
2750
2755
2760
2765
2770
2775
2780
2785
2790
2795
2800
2805
2810
2815
2820
2825
2830
2835
2840
2845
2850
2855
2860
2865
2870
2875
2880
2885
2890
2895
2900
2905
2910
2915
2920
2925
2930
2935
2940
2945
2950
2955
2960
2965
2970
2975
2980
2985
2990
2995
3000
3005
3010
3015
3020
3025
3030
3035
3040
3045
3050
3055
3060
3065
3070
3075
3080
3085
3090
3095
3100
3105
3110
3115
3120
3125
3130
3135
3140
3145
3150
3155
3160
3165
3170
3175
3180
3185
3190
3195
3200
3205
3210
3215
3220
3225
3230
3235
3240
3245
3250
3255
3260
3265
3270
3275
3280
3285
3290
3295
3300
3305
3310
3315
3320
3325
3330
3335
3340
3345
3350
3355
3360
3365
3370
3375
3380
3385
3390
3395
3400
3405
3410
3415
3420
3425
3430
3435
3440
3445
3450
3455
3460
3465
3470
3475
3480
3485
3490
3495
3500
3505
3510
3515
3520
3525
3530
3535
3540
3545
3550
3555
3560
3565
3570
3575
3580
3585
3590
3595
3600
3605
3610
3615
3620
3625
3630
3635
3640
3645
3650
3655
3660
3665
3670
3675
3680
3685
3690
3695
3700
3705
3710
3715
3720
3725
3730
3735
3740
3745
3750
3755
3760
3765
3770
3775
3780
3785
3790
3795
3800
3805
3810
3815
3820
3825
3830
3835
3840
3845
3850
3855
3860
3865
3870
3875
3880
3885
3890
3895
3900
3905
3910
3915
3920
3925
3930
3935
3940
3945
3950
3955
3960
3965
3970
3975
3980
3985
3990
3995
4000
4005
4010
4015
4020
4025
4030
4035
4040
4045
4050
4055
4060
4065
4070
4075
4080
4085
4090
4095
4100
4105
4110
4115
4120
4125
4130
4135
4140
4145
4150
4155
4160
4165
4170
4175
4180
4185
4190
4195
4200
4205
4210
4215
4220
4225
4230
4235
4240
4245
4250
4255
4260
4265
4270
4275
4280
4285
4290
4295
4300
4305
4310
4315
4320
4325
4330
4335
4340
4345
4350
4355
4360
4365
4370
4375
4380
4385
4390
4395
4400
4405
4410
4415
4420
4425
4430
4435
4440
4445
4450
4455
4460
4465
4470
4475
4480
4485
4490
4495
4500
4505
4510
4515
4520
4525
4530
4535
4540
4545
4550
4555
4560
4565
4570
4575
4580
4585
4590
4595
4600
4605
4610
4615
4620
4625
4630
4635
4640
4645
4650
4655
4660
4665
4670
4675
4680
4685
4690
4695
4700
4705
4710
4715
4720
4725
4730
4735
4740
4745
4750
4755
4760
4765
4770
4775
4780
4785
4790
4795
4800
4805
4810
4815
4820
4825
4830
4835
4840
4845
4850
4855
4860
4865
4870
4875
4880
4885
4890
4895
4900
4905
4910
4915
4920
4925
4930
4935
4940
4945
4950
4955
4960
4965
4970
4975
4980
4985
4990
4995
5000
5005
5010
5015
5020
5025
5030
5035
5040
5045
5050
5055
5060
5065
5070
5075
5080
5085
5090
5095
5100
5105
5110
5115
5120
5125
5130
5135
5140
5145
5150
5155
5160
5165
5170
5175
5180
5185
5190
5195
5200
5205
5210
5215
5220
5225
5230
5235
5240
5245
5250
5255
5260
5265
5270
5275
5280
5285
5290
5295
5300
5305
5310
5315
5320
5325
5330
5335
5340
5345
5350
5355
5360
5365
5370
5375
5380
5385
5390
5395
5400
5405
5410
5415
5420
5425
5430
5435
5440
5445
5450
5455
5460
5465
5470
5475
5480
5485
5490
5495
5500
5505
5510
5515
5520
5525
5530
5535
5540
5545
5550
5555
5560
5565
5570
5575
5580
5585
5590
5595
5600
5605
5610
5615
5620
5625
5630
5635
5640
5645
5650
5655
5660
5665
5670
5675
5680
5685
5690
5695
5700
5705
5710
5715
5720
5725
5730
5735
5740
5745
5750
5755
5760
5765
5770
5775
5780
5785
5790
5795
5800
5805
5810
5815
5820
5825
5830
5835
5840
5845
5850
5855
5860
5865
5870
5875
5880
5885
5890
5895
5900
5905
5910
5915
5920
5925
5930
5935
5940
5945
5950
5955
5960
5965
5970
5975
5980
5985
5990
5995
6000
6005
6010
6015
6020
6025
6030
6035
6040
6045
6050
6055
6060
6065
6070
6075
6080
6085
6090
6095
6100
6105
6110
6115
6120
6125
6130
6135
6140
6145
6150
6155
6160
6165
6170
6175
6180
6185
6190
6195
6200
6205
6210
6215
6220
6225
6230
6235
6240
6245
6250
6255
6260
6265
6270
6275
6280
6285
6290
6295
6300
6305
6310
6315
6320
6325
6330
6335
6340
6345
6350
6355
6360
6365
6370
6375
6380
6385
6390
6395
6400
6405
6410
6415
6420
6425
6430
6435
6440
6445
6450
6455
6460
6465
6470
6475
6480
6485
6490
6495
6500
6505
6510
6515
6520
6525
6530
6535
6540
6545
6550
6555
6560
6565
6570
6575
6580
6585
6590
6595
6600
6605
6610
6615
6620
6625
6630
6635
6640
6645
6650
6655
6660
6665
6670
6675
6680
6685
6690
6695
6700
6705
6710
6715
6720
6725
6730
6735
6740
6745
6750
6755
6760
6765
6770
6775
6780
6785
6790
6795
6800
6805
6810
6815
6820
6825
6830
6835
6840
6845
6850
6855
6860
6865
6870
6875
6880
6885
6890
6895
6900
6905
6910
6915
6920
6925
6930
6935
6940
6945
6950
6955
6960
6965
6970
6975
6980
6985
6990
6995
7000
7005
7010
7015
7020
7025
7030
7035
7040
7045
7050
7055
7060
7065
7070
7075
7080
7085
7090
7095
7100
7105
7110
7115
7120
7125
7130
7135
7140
7145
7150
7155
7160
7165
7170
7175
7180
7185
7190
7195
7200
7205
7210
7215
7220
7225
7230
7235
7240
7245
7250
7255
7260
7265
7270
7275
7280
7285
7290
7295
7300
7305
7310
7315
7320
7325
7330
7335
7340
7345
7350
7355
7360
7365
7370
7375
7380
7385
7390
7395
7400
7405
7410
7415
7420
7425
7430
7435
7440
7445
7450
7455
7460
7465
7470
7475
7480
7485
7490
7495
7500
7505
7510
7515
7520
7525
7530
7535
7540
7545
7550
7555
7560
7565
7570
7575
7580
7585
7590
7595
7600
7605
7610
7615
7620
7625
7630
7635
7640
7645
7650
7655
7660
7665
7670
7675
7680
7685
7690
7695
7700
7705
7710
7715
7720
7725
7730
7735
7740
7745
7750
7755
7760
7765
7770
7775
7780
7785
7790
7795
7800
7805
7810
7815
7820
7825
7830
7835
7840
7845
7850
7855
7860
7865
7870
7875
7880
7885
7890
7895
7900
7905
7910
7915
7920
7925
7930
7935
7940
7945
7950
7955
7960
7965
7970
7975
7980
7985
7990
7995
8000
8005
8010
8015
8020
8025
8030
8035
8040
8045
8050
8055
8060
8065
8070
8075
8080
8085
8090
8095
8100
8105
8110
8115
8120
8125
8130
8135
8140
8145
8150
8155
8160
8165
8170
8175
8180
8185
8190
8195
8200
8205
8210
8215
8220
8225
8230
8235
8240
8245
8250
8255
8260
8265
8270
8275
8280
8285
8290
8295
8300
8305
8310
8315
8320
8325
8330
8335
8340
8345
8350
8355
8360
8365
8370
8375
8380
8385
8390
8395
8400
8405
8410
8415
8420
8425
8430
8435
8440
8445
8450
8455
8460
8465
8470
8475
8480
8485
8490
8495
8500
8505
8510
8515
8520
8525
8530
8535
8540
8545
8550
8555
8560
8565
8570
8575
8580
8585
8590
8595
8600
8605
8610
8615
8620
8625
8630
8635
8640
8645
8650
8655
8660
8665
8670
8675
8680
8685
8690
8695
8700
8705
8710
8715
8720
8725
8730
8735
8740
8745
8750
8755
8760
8765
8770
8775
8780
8785
8790
8795
8800
8805
8810
8815
8820
8825
8830
8835
8840
8845
8850
8855
8860
8865
8870
8875
8880
8885
8890
8895
8900
8905
8910
8915
8920
8925
8930
8935
8940
8945
8950
8955
8960
8965
8970
8975
8980
8985
8990
8995
9000
9005
9010
9015
9020
9025
9030
9035
9040
9045
9050
9055
9060
9065
9070
9075
9080
9085
9090
9095
9100
9105
9110
9115
9120
9125
9130
9135
9140
9145
9150
9155
9160
9165
9170
9175
9180
9185
9190
9195
9200
9205
9210
9215
9220
9225
9230
9235
9240
9245
9250
9255
9260
9265
9270
9275
9280
9285
9290
9295
9300
9305
9310
9315
9320
9325
9330
9335
9340
9345
9350
9355
9360
9365
9370
9375
9380
9385
9390
9395
9400
9405
9410
9415
9420
9425
9430
9435
9440
9445
9450
9455
9460
9465
9470
9475
9480
9485
9490
9495
9500
9505
9510
9515
9520
9525
9530
9535
9540
9545
9550
9555
9560
9565
9570
9575
9580
9585
9590
9595
9600
9605
9610
9615
9620
9625
9630
9635
9640
9645
9650
9655
9660
9665
9670
9675
9680
9685
9690
9695
9700
9705
9710
9715
9720
9725
9730
9735
9740
9745
9750
9755
9760
9765
9770
9775
9780
9785
9790
9795
9800
9805
9810
9815
9820
9825
9830
9835
9840
9845
9850
9855
9860
9865
9870
9875
9880
9885
9890
9895
9900
9905
9910
9915
9920
9925
9930
9935
9940
9945
9950
9955
9960
9965
9970
9975
9980
9985
9990
9995
10000
10005
10010
10015
10020
10025
10030
10035
10040
10045
10050
10055
10060
10065
10070
10075
10080
10085
10090
10095
10100
10105
10110
10115
10120
10125
10130
10135
10140
10145
10150
10155
10160
10165
10170
10175
10180
1

Corey et al Tetrahedron Lett 1976, 3-6; and Corey et al., Tetrahedron Lett, 1976, 3667-3668]. The modifications are provided in Example 1 below. Initial screening of epimeric 16-ethyl-2-methoxyestradiol and related analogues showed that it is about equipotent to 2-methoxyestradiol in inhibition of HUVEC cell proliferation *in vitro*.

Administration

The compositions described above can be provided as physiologically acceptable formulations using known techniques, and these formulations can be administered by standard routes. In general, the combinations may be administered by the topical, oral, rectal or parenteral (e.g., intravenous, subcutaneous or intramuscular) route. In addition, the combinations may be incorporated into biodegradable polymers allowing for sustained release, the polymers being implanted in the vicinity of where delivery is desired, for example, at the site of a tumor. The biodegradable polymers and their use are described in detail in Brem et al., *J. Neurosurg.* 74:441-446 (1991). The dosage of the composition will depend on the condition being treated, the particular derivative used, and other clinical factors such as weight and condition of the patient and the route of administration of the compound. However, for oral administration to humans, a dosage of 0.01 to 100 mg/kg/day, preferably 0.01-1 mg/kg/day, is generally sufficient.

The formulations include those suitable for oral, rectal, nasal, topical (including buccal and sublingual), vaginal or parenteral (including subcutaneous, intramuscular, intravenous, intradermal, intraocular, intratracheal, and epidural) administration. The formulations may conveniently be presented in unit dosage form and may be prepared by conventional pharmaceutical techniques. Such techniques include the step of bringing into association the active ingredient and the pharmaceutical carrier(s) or excipient(s). In general, the formulations are prepared by uniformly and intimately bringing into association the active ingredient with liquid carriers or finely divided solid carriers or both, and then, if necessary, shaping the product.

Formulations of the present invention suitable for oral administration may be presented as discrete units such as capsules,

5 cachets or tablets each containing a predetermined amount of the active ingredient; as a powder or granules; as a solution or a suspension in an aqueous liquid or a non-aqueous liquid; or as an oil-in-water liquid emulsion or a water-in-oil emulsion and as a bolus, etc.

10 A tablet may be made by compression or molding, optionally with one or more accessory ingredients. Compressed tablets may be prepared by compressing, in a suitable machine, the active ingredient in a free-flowing form such as a powder or granules, optionally mixed with a binder, lubricant, inert diluent, preservative, surface-active or dispersing agent. Molded tables may be made by molding, in a suitable machine, a mixture of the powdered compound moistened with an inert liquid diluent. The tablets may optionally coated or scored and may be formulated so as to provide a slow or controlled release of the active ingredient therein.

15 Formulations suitable for topical administration in the mouth include lozenges comprising the ingredients in a flavored basis, usually sucrose and acacia or tragacanth; pastilles comprising the active ingredient in an inert basis such as gelatin and glycerin, or sucrose and acacia; and mouthwashes comprising the ingredient to be administered in a suitable liquid carrier.

20 Formulations suitable for topical administration to the skin may be presented as ointments, creams, gels and pastes comprising the ingredient to be administered in a pharmaceutical acceptable carrier. A preferred topical delivery system is a transdermal patch containing the ingredient to be administered.

25 Formulations for rectal administration may be presented as a suppository with a suitable base comprising, for example, cocoa butter or a salicylate.

30 Formulations suitable for nasal administration, wherein the carrier is a solid, include a coarse powder having a particle size, for example, in the range of 20 to 500 microns which is administered in the manner in which snuff is taken, i.e., by rapid inhalation through the nasal passage from a container of the powder held close up to the nose. Suitable formulations, wherein the carrier is a liquid, for administration, as for example, a nasal spray or as nasal drops, include aqueous or oily solutions of the active ingredient.

Formulations suitable for vaginal administration may be presented as pessaries, tampons, creams, gels, pastes, foams or spray formulations containing in addition to the active ingredient such as carriers as are known in the art to be appropriate.

Formulations suitable for parenteral administration include aqueous and non-aqueous sterile injection solutions which may contain anti-oxidants, buffers, bacteriostats and solutes which render the formulation isotonic with the blood of the intended recipient; and aqueous and non-aqueous sterile suspensions which may include suspending agents and thickening agents. The formulations may be presented in unit-dose or multi-dose containers, for example, sealed ampules and vials, and may be stored in a freeze-dried (lyophilized) conditions requiring only the addition of the sterile liquid carrier, for example, water for injections, immediately prior to use. Extemporaneous injection solutions and suspensions may be prepared from sterile powders, granules and tables of the kind previously described.

Preferred unit dosage formulations are those containing a daily dose or unit, daily sub-dose, as herein above recited, or an appropriate fraction thereof, of the administered ingredient.

2-Methoxyestradiol is an endogenous metabolite of estradiol (E2) that has potent anti-proliferative activity and induces apoptosis in a wide variety of tumor and non-tumor cell lines. When administered orally, it exhibits anti-tumor and anti-proliferative activity with little or no toxicity. *In vitro* data suggests that 2-methoxyestradiol does not engage the estrogen receptor for its anti-proliferative activity and is not estrogenic over a wide range of concentrations, as assessed by estrogen dependant MCF-7 cell proliferation. However, the presence of demethylases *in vivo* may metabolize this compound to 2-hydroxyestradiol, which has been shown to be estrogenic by several approaches. The present invention improves the bioavailability of estradiol or 2-methoxyestradiol and to reduces the formation of estrogenic 2-methoxyestradiol metabolites. The present invention modifies estradiol or 2-methoxyestradiol in such a way that the molecule can not be converted into an uterotrophic derivative.

One embodiment of the invention modifies the methyl ether of 2-methoxyestradiol so that it can not be a substrate for

demethylase. Additionally, it has been demonstrated (Cushman et al *J. Med. Chem.* **1995**, 38, 2041-2049) that other electron-rich groups at the 2-position of estradiol (propyne, propene, ethoxy) have good anti-proliferative activity *in vitro*. It is disclosed that modifications at C-2 of estradiol such as formyl, acetyl, methanol, 1-ethanol, 2-ethanol, amino, alkylamino, dialkyl amino, methyleneamine, methylene alkyl amine and methylene dialkylamine, and alkyl amide are be anti-proliferative and anti-angiogenic agents have reduced or removed uterotrophic activity. Alkyl is defined as any carbon chain up to 6 carbons in length that is branched or straight. Listed below in Table 1 are data of 2-modified estradiol derivatives in HUVEC, MDA-MB-231 and MCF7 proliferation data. The synthetic paths for preparation of these analogs can be found in Pert et al *Aust. J. Chem.* **1989**, 42, 405-419. Lovely et al *Tetrahedron Lett.* **1994**, 35, 8735-8738. Gonzalez et al *Steroids* **1982**, 40, 171-187. Nambara et al *Chem. Pharm. Bull.* **1970**, 18, 474-480. Cushman et al *J. Med. Chem.* **1995**, 38, 2041-2049 and methods developed in-house and are discussed below.

Table 1

Compound	HUVEC (IC ₅₀ μM)	MDA-MB-231 (IC ₅₀ μM)	MCF7 Proliferation Index
E2	NA	NA	13.1
2ME2	0.5	0.9	4.4
2-methyl hydroxy-E2	10	>25	7.4
2-formyl-E2	8	>25	5.4
2-acetyl-E2	18	9	4.4

All of the 2-modified analogs presented in Table 1 have significantly less estrogenic activity (compared to estradiol) as represented by their proliferation index in estrogen dependant MCF-7 cells. All of these analogs have the capacity to form a hydrogen bond with the hydroxy group at position 3 and this may be the reason for their relatively low estrogenic character compared to estradiol. Both the 2-methylhydroxy and 2-formyl derivatives had good antiproliferative activity (IC₅₀ < 10 microM) in HUVEC cells, whereas the 2-acetyl had poor activity in the same assay. In contrast,


2-methylhydroxy and 2-formyl were inactive in breast tumor MDA-MB-231 cells while 2-acetyl E2 had good activity in this cell line.

Although not wishing to be bound by theory, molecular modeling suggests that there may be a hydrogen bond that forms between the 3-hydroxy group and the methoxy group of 2-methoxyestradiol. This interaction may be important for both 2-methoxyestradiol's anti-proliferative and anti-angiogenic action as well as its non-estrogenic activity. It is claimed that any group that can be placed at position 2 of estradiol and has the potential to form a hydrogen bond with the 3-hydroxy group is an anti-proliferative and anti-angiogenic agent that lacks estrogenic activity.

It should be understood that in addition to the ingredients, particularly mentioned above, the formulations of this invention may include other agents convention in the art having regard to the type of formulation in question, for example, those suitable for oral administration may include flavoring agents.

EXPERIMENTAL DATA

The following Examples refer to the compound of the general formula:



wherein:

a) R_b and R_o are independently -H, -Cl, -Br, -I, -F, -CN, lower alkyl, -OH, -CH₂-OH, -NH₂; or N(R₆)(R₇), wherein R₆ and R₇ are independently hydrogen or an alkyl or branched alkyl with up to 6 carbons;

~~Sub A2~~
b) R_a is $-N_3$, $-C \equiv N$, $-C \equiv C-R$, $-C=CH-R$, $-R-C=CH_2$, $-C \equiv CH$, $-O-R$, $-R-R_1$, or $-O-R-R_1$ where R is a straight or branched alkyl with up to 10 carbons or aralkyl, and R_1 is $-OH$, $-NH_2$, $-Cl$, $-Br$, $-I$, $-F$ or CF_3 ;

5 c) Z' is $>CH$, $>COH$, or $>C-R_2-OH$, where R_2 is an alkyl or branched alkyl with up to 10 carbons or aralkyl;

10 d) $>C-R_g$ is $>CH_2$, $>C(H)-OH$, $>C=O$, $>C=N-OH$, $>C(R_3)OH$, $>C=N-OR_3$, $>C(H)-NH_2$, $>C(H)-NHR_3$, $>C(H)-NR_3R_4$, or $>C(H)-C(O)-R_3$, where each R_3 and R_4 is independently an alkyl or branched alkyl with up to 10 carbons or aralkyl;

~~Sub A3~~
15 e) R_{h1} and R_{h2} are independently H , or a straight or branched chain alkyl, alkenyl or alkynyl with up to 6 carbons that is unsubstituted, or substituted with one or more groups selected from a hetero functionality ($O-Y$, $N-Y$ or $S-Y$) where Y is H , Me or an alkyl chain up to 6 carbons; a halo functionality (F , Cl , Br or I); an aromatic group optionally substituted with hetero, halo or alkyl; or R_{h1} and R_{h2} are independently an aromatic group optionally substituted with hetero, halo or alkyl, provided that both R_{h1} and R_{h2} are not H ;

~~Sub A4~~
25 f) Z'' is $>CH_2$, $>C=O$, $>C(H)-OH$, $>C=N-OH$, $>C=N-OR_5$, $>C(H)-C \equiv N$, or $>C(H)-NR_5R_5$, wherein each R_5 is independently hydrogen, an alkyl or branched alkyl with up to 10 carbons or aralkyl;

30 and wherein all monosubstituted substituents have either an α or β configuration.

Lower alkyl is defined as a small carbon chain having 1-8 carbon atoms. The chain may be branched or unbranched.

EXAMPLE 1

Synthesis of 2-ME Derivatives and modifications at the 16 position

Synthesis of the 2-ME derivatives described herein is within the capability of one ordinarily skilled in the art. A specific description of the synthesis of the 2-ME derivatives having modifications at the 2 and 6 positions and analogs discussed herein can be found in M. Cushman, H-M. He, J.A. Katzenellenbogen, C.M. Lin and E. Hamel, Synthesis, antitubulin and antimitotic activity, and cytotoxicity of 2-methoxyestradiol, and endogenous mammalian metabolite of estradiol that inhibits tubulin polymerization by binding to the colchicine binding site, *J. Med. Chem.*, 38(12): 2042 (1995); and M. Cushman, H-M. He, J. Katzenellenbogen, R. Varma, E. Hamel, C. Lin, S. Ram and Y.P. Sachdeva, Synthesis of analogs of 2-methoxyestradiol with enhanced inhibitory effects on tubulin polymerization and cancer cell growth, *J. Med. Chem.* 40(15): 2323 (1997).

The synthetic pathways used to prepare the derivatives of the estradiol derivatives modified at the 16 position of the present invention are based on modified published literature procedures for estradiol derivatives cited earlier. Examples of the modifications are provided in Examples 2 through 23 below.

EXAMPLE 2

Preparation of 3-Benzyl-2-methoxyestradiol

2-Methoxyestradiol (10.09 g, 33.4 mmol) and potassium carbonate (22 g, 278 mmol) were suspended in anhydrous ethanol and cooled to 0°C. Benzyl bromide (11.4 mL, 95.8 mmol) was added dropwise, and following the addition, the mixture was brought to reflux for 8 h. The solution was cooled to room temperature (rt), and the solvent was removed via rotoevap. The resulting residue was diluted with approximately 200 ml water, and washed with ethyl acetate (3 x 200 mL). The combined organics were washed with water (200 mL), sodium bicarbonate (saturated (satd), 200mL) and brine (200 mL). Dry with sodium sulfate, filter and roto-evaporation (rotoevap). Product was dried under vacuo with occasional gentle heating using a heat gun to give a yellowish glass (13.54 g, quantitative yield) and used without further purification.

Selected spectral data: $^1\text{H-NMR}$ (300 MHz, CDCl_3) δ 7.29-7.53 (m, 5H), 6.88 (s, 1H), 6.65 (s, 3H), 5.11 (s, 2H), 3.87 (s, 3H), 3.7 (t, $J=8$ Hz, 1H), 0.80 (s, 3H). FT-IR (neat) 3341, 2920, 2864, 1605, 1513, 1453, 1254, 1211, 1117, 1022 cm^{-1} .

EXAMPLE 3

Preparation of 3-Benzyl-2-methoxyestrone

Oxalyl chloride (38 mmol, 19 mL, 2M, methylene chloride) was added to anhydrous methylene chloride (25 mL) and cooled to -46°C . Methyl sulfoxide (5.40 mL, 76 mmol) was added dropwise, and the mixture was stirred for 2 minutes. 3-Benzyl-2-methoxyestradiol in methylene chloride/methyl sulfoxide (10 mL/15 mL) and added within 5 minutes and the resulting mixture was stirred for 1 h. Triethyl amine (170 mmol, 23.5 mL) was added drop-wise, stirred 5 minutes and warmed to rt. Water (~200 mL) was added and the mixture was washed with methylene chloride (3x 200 mL). The combined organics were washed with water (200 mL), dilute HCl (1% aq., 200 mL), sodium carbonate (satd, 200 mL) and brine (200 mL). The organics were dried with magnesium sulfate, filtered and rotoevaped to give a white solid. The solid was crystallized with hot ethanol to give white crystals (9.94g, 25.5 mmol, 76% overall yield from 2-methoxyestradiol).

Selected spectral data: $^1\text{H-NMR}$ (300 MHz, CDCl_3) δ 7.28-7.48 (m, 5H), 6.86 (s, 1H), 6.66 (s, 1H), 3.88 (s, 3H), 0.94 (s, 3H). IR (neat) 2920, 1731, 1519, 1202, 1012 cm^{-1} .

EXAMPLE 4

Representative preparation of 16 α -alkyl-3-benzyl-2-methoxyestrone

Lithium diisopropyl amide (2M, Aldrich, heptane/THF/ethylbenzene) was dissolved in THF and cooled to -78°C , and 3-benzyl-2-methoxyestrone in THF (10 mL) was added dropwise. Following addition, the mixture was warmed to 0°C and stirred 1 hour (h). The mixture was then cooled to -78°C and DMPU (1mL) followed by crotyl bromide (205 μL , 2.0 mmol) were added dropwise. The mixture was warmed to rt over 4 h. The reaction was quenched by carefully adding water (100 mL) and

washing with ethyl acetate (2 x 100 mL). The combined organics were washed with water (100 mL) and brine (100 mL). The solution was dried with magnesium sulfate, filtered and rotoevaped. The crude product was purified using hexane / ethyl acetate (9:1) SiO₂ Biotage FLASH apparatus. 680 mg (1.53 mmol) of product was obtained and approximately 121 mg (0.31 mmol) of starting material was recovered (90% yield based on recovered starting material). Diastereomeric ratio of 16 α / β is approximately 2:1 (s H18 signals at 0.88, 0.79 ppm).

Selected spectral data: ¹H-NMR (300 MHz, CDCl₃) δ 7.28-7.48 (m, 5H), 6.86 (s, 1H), 6.66 (s, 1H), 5.34-5.59 (m, 2H), 5.13 (s, 2H), 3.88 (s, 3H), 0.87 & 0.97 (s, total 3H, ratio 1:2).

EXAMPLE 5

Representative preparation of 16 β -alkyl-3-benzyl-2-methoxyestrone

3-Benzyl-2-methoxyestrone (1.175g, 3.0 mmol) was dissolved in anhydrous THF (15 mL), cooled to -78°C and lithium diisopropyl amide (2M Aldrich, heptane/THF/ethylbenzene) was added dropwise and stirred 1h. DMPU (1 mL) followed by crotyl bromide (302 μ L) were added and the mixture warmed to rt over 24 h. Workup as above and purify using hexane:ethyl acetate (4:1) SiO₂ flash column gave 492 mg purified product (1.1 mol, 37% yield).

Selected spectral data: ¹H-NMR (300 MHz, CDCl₃) δ 7.28-7.48 (m, 5H), 6.86 (s, 1H), 6.66 (s, 1H), 5.62-5.34 (m, 2H), 5.13 (s, 2H), 3.89 (s, 3H), 0.98 and 0.87 (s, 3H total, ratio 2:1). IR (neat) 2928, 2854, 1732, 1606, 1508, 1452, 1215, 1016 cm⁻¹.

EXAMPLE 6

Representative preparation of 16 β -alkyl-3-benzyl-2-methoxyestrone

3-benzyl-16-carbomethoxy-2-methoxyestrone (0.840 g, 1.87 mmol), potassium hydride (1.5 g, 10.9 mmol, 30% mineral oil dispersion, washed in hexanes) and 18-crown-6 (120 mg, 0.4 mmol) was mixed in THF (40 mL) and refluxed for 1 h. The mixture was cooled to rt, and allyl bromide (537 μ L, 6.2 mmol) was added and the mixture was refluxed for 18 h. After cooling to rt, the reaction was quenched by carefully adding approximately 2 ml of water with

stirring, then adding an additional 100 mL water. This mixture was washed with ethyl acetate (2 x 100 mL) and the combined organics were washed with brine (100 mL). The organics were dried with magnesium sulfate, filtered and rotoevaped. Purification using 85:5 hexanes:ethyl acetate SiO₂ Biotage FLASH apparatus yielded 697 mg of product (1.42 mol, 76% yield).

Selected spectral data: ¹H-NMR (300 MHz, CDCl₃) δ 7.28-7.48 (m, 5H), 6.85 (s, 1H), 6.66 (s, 1H), 5.66-5.79 (m, 1H), 5.15-5.20 (m, 2H), 5.13 (s, 2H), 3.88 (s, 3H), 3.75 (s, 3H), 0.99 (s, 3H).

EXAMPLE 7

Representative decarboxylation of 16-alkyl-16-carbomethoxy-3-benzyl-2-methoxyestrone

16-allyl-16-carbomethoxy-3-benzyl-2-methoxyestrone (697 mg, 1.42 mmol), lithium chloride (1.15 g, 27 mmol), water (485 μL, 27 mmol) were dissolved in DMF (63 mL) and refluxed for 20 h. Cool to rt, add 1N HCl (100 mL) and wash with ether (2 x 100 mL) the combined organics were washed with water (100 mL), and brine 100 mL), dry with magnesium sulfate, filter and rotoevap. Purification by 85:15 hexanes:ethyl acetate SiO₂ Biotage Flash apparatus gave 271 mg product and 189 mg recovered starting material. Starting material was resubjected to the reaction (308 mg LiCl, 132 μL, water, 17 mL DMF) for 28 h and worked up as above to give 130 mg product. Overall yield for reaction was 66% (401 mg, 0.93 mmol).

Selected spectral data: ¹H-NMR (300 MHz, CDCl₃) δ 7.28-7.48 (m, 5H), 6.85 (s, 1H), 6.65 (s, 1H), 5.69-5.88 (m, 1H), 5.13 (s, 2H), 5.00-5.08 (m, 2H), 5.88 (s, 3H), 0.98 nd 0.88 (s, total 3H, ratio 1:1.4). FT-IR (neat), 2925, 2855, 1726, 1514, 1214, 1103 cm⁻¹.

EXAMPLE 8

Preparation of 16-methane-dimethylenamine-3-benzyl-2-methoxyestrone

3-benzyl-2-methoxyestrone (1.51 g, 3.87 mmol) was suspended in *tert*-butoxy bis(dimethylamino)methane (1.64 mL, 8.13 mmol) and heated in an oil bath (155°C) for 1.5 h, during which time

the steroid dissolved. The reaction mixture was cooled to rt, and poured into ice water (100 mL) and washed with methylene chloride (2 x 100 mL). The organics were washed with brine (100 mL) dried with magnesium sulfate, filtered and rotoevaped to give product which was used without further purification (1.82 g, quantitative yield).

Selected spectral data: $^1\text{H-NMR}$ (300 MHz, CDCl_3) δ 7.23-7.47 (m, 5H), 6.87 (s, 1H), 6.64 (s, 1H), 5.12 (s, 2H), 3.88 (s, 3H), 3.07 (s, 6H), 0.91 (s, 3H).

EXAMPLE 9

Preparation of 16-carbomethoxy-3-benzyl-2-methoxy estrone

3-Benzyl-2-methoxyestrone (1.6113g, 2.978 mmol) was dissolved in THF (15 mL), cooled to -78°C and lithium diisopropyl amide (2M, Aldrich, Heptane / THF / ethylbenzene) was added dropwise and stirred for 1 h. Methyl cyanoformate (237 μL , 3 mmol) in DMPU (1mL) was added and the mixture warmed to rt over 18 h. Water (100 ml) was carefully added, and the mixture was washed with ethyl acetate (3 x 100 mL) and the combined organics were washed with brine (100 mL), dried with sodium sulfate, filtered and rotoevaped. Final purification of product using hexane:ethyl acetate (85:15) then switching to hexane: ethyl acetate (75:25) SiO_2 flash column yielded 806 mg product (1.8 mmol, 60%).

Selected spectral data: $^1\text{H-NMR}$ (300 MHz, CDCl_3) δ 7.28-7.48 (m, 5H), 6.85 (s, 1H), 6.66 (s, 1H), 5.13 (s, 2H), 3.88 (s, 3H), 3.78 (s, 3H), 3.23 (dd, $J = 9, 10$ Hz, 1H), 1.0 (s, 3H). FT-IR (neat) 2929, 2860, 1750, 1723, 1604, 1508, 1211, 1014 cm^{-1} .

EXAMPLE 10

Representative procedure for preparation of 16-alkyl-3-benzyl-2-methoxyestra-17 β -diol

16 α -crotyl-3-benzyl-2methoxyestrone (680 mg, 1.53 mmol) was dissolved in anhydrous THF (10 mL), and cooled to -78°C . Lithium aluminum hydride (3.06 mmol, 116 mg) was added and the solution was stirred for 2 h. The reaction was quenched by carefully adding water (2 mL) and warming to rt, then adding additional 50 mL portion of water. The mixture was washed with ethyl acetate (2 x 50 mL) and the combined organics were washed

with water (50 mL), brine (50 mL), dried with magnesium sulfate, filtered and rotoevaped. The mixture was purified with 3:1 hexane:ethyl acetate SiO₂ Biotage FLASH apparatus to give 500 mg purified product (1.12 mmol, 73% yield).

Selected spectral data: ¹H-NMR (300 MHz, CDCl₃) δ 7.28-7.48 (m, 5H), 6.87 (s, 1H), 6.64 (s, 1H), 5.47-5.56 (m, 2H), 5.12 (s, 2H), 3.88 (s, 3H), 3.8 (d, J=9 Hz) and 3.33 (d, J=8Hz) total 1H, ratio 1:1.7, 0.84 and 0.81 (s, 3H total).

EXAMPLE 11

Preparation of 16-methanol-3-benzyl-2-methoxyestradiol

Reaction procedure and work up as above, (used 806 mg, 1.8 mmol 16-carbomethoxy-3-benzyl-2-methoxyestrone), except warm to rt for 2 h before quenching. Purify final product with 3:2 hexane:ethyl acetate SiO₂ flash column. Obtain 304 mg β isomer, 51 mg α isomer which were separated by chromatography. Selected spectral data: ¹H-NMR (300 MHz, CDCl₃) δ Major isomer 7.28-7.48 (m, 5H), 6.87 (s, 1H), 6.64 (s, 1H), 5.12 (s, 2H), 3.97 (d, J=10 Hz), 3.88 (s and obscured d, 4H), 3.67 (dd, J=4, 7Hz, 1H), 0.87 (s, 3H). Minor isomer 7.28-7.47 (m, 5H), 6.86 (s, 1H), 6.64 (s, 1H), 3.88 (s, 3H), 3.83 (d, J=14Hz, 1H), 3.69 (t, J=9Hz, 1H), 3.54 (d, J=7Hz, 1H), 0.87 (s, 3H).

EXAMPLE 12

Representative debenzylation of 16-alkyl-3-benzyl-2-methoxyestradiol

16α-crotyl-3-benzyl-2-methoxyestradiol (500 mg, 1.12 mmol) was dissolved in ethyl acetate (25 mL) in Parr reaction bottle. The bottle was flushed with argon, and Pd /C (10%, 2.5 g) was added. The bottle was fitted to a Parr hydrogenator, filled and purged with hydrogen five times, pressurized to 50 psi, and agitated for 24 h. The mixture was filtered through a celite pad, rotoevaped and purified with a 3:1 hexane ethyl acetate SiO₂ flash column. Obtain 358 mg product (1.0 mmol, 89%).

Selected spectral data: ¹H-NMR (300 MHz, CDCl₃) δ 6.81 (s, 1H), 6.66 (s, 1H), 3.87 (s, 3H), 3.76 (d, J=10 Hz) and 3.29 (d, J=8Hz) (total 1H, ratio 1:2), 0.82 and 0.79 (s, 3H). FT-IR (neat) 3245, 2914, 1606, 1523, 1414, 1258, 1028 cm⁻¹. Analysis calculated

(Anal. Calcd) for $C_{20}H_{34}O_3$: C, 77.44; H, 9.56. Found: C, 76.64; H, 9.51.

EXAMPLE 13

16 β -methyl-2methoxyestradiol

Selected spectral data: 1H -NMR (300 MHz, $CDCl_3$) δ 6.81 (s, 1H), 6.66 (s, 1H), 3.87 (s, 3H), 3.73 (d, J=10 Hz) and 3.23 (d, J=8 Hz) (total 1H, 2:1), 0.83 and 0.81 (s, 3 H total). Anal. Calcd for $C_{20}H_{28}O_3$, 1/4 H_2O : C, 74.85; H, 8.95. Found: C, 74.93; H, 8.94.

EXAMPLE 14

16 α -methyl-2methoxyestradiol

Selected spectral data: 1H -NMR (300 MHz, $CDCl_3$) δ 6.81 (s, 1H), 6.66 (s, 1H), 3.87 (s, 3H), 3.23 (d, J=7 Hz) (s, 1H), 0.81 (s, 3 H). Anal. Calcd for $C_{20}H_{28}O_3$, 1/4 H_2O : C, 74.85; H, 8.95. Found: C, 74.98; H, 8.65.

EXAMPLE 15

Racemic 16-ethyl-2-methoxyestradiol

Selected spectral data: 1H -NMR (300 MHz, $CDCl_3$) δ 6.82 (s, 1H), 6.66 (s, 1H), 3.88 (s, 3H), 3.76 (d, J=9 Hz) and 3.30 (d, J=10 Hz), (1H total, ratio 1:1), 0.83 and 0.79 (s, 3H total). FT-IR (neat) 3214, 2918, 1605, 1522, 1229, 1201, 1024 cm^{-1} . Anal. Calcd for $C_{21}H_{30}O_3$: C, 76.33; H, 9.15. Found: C, 76.18; H, 9.16.

EXAMPLE 16

16 α -n-propyl-2-methoxyestradiol

Selected spectral data: 1H -NMR (300 MHz, $CDCl_3$) δ 6.81 (s, 1H), 6.66 (s, 1H), 5.43 (s, 1H), 3.87 (s, 3H), 3.29 (t, J=7 Hz, 1H), 0.95 (t, J=7 Hz, 3H), 0.83 and 0.80 (s, total 3H, ratio 7.3:1). Anal. Calcd for $C_{22}H_{32}O_3$: C, 76.69; H, 9.37. Found: C, 76.55; H, 9.44.

EXAMPLE 17

16 β -n-propyl-2-methoxyestradiol

Selected spectral data: ¹H-NMR (300 MHz, CDCl₃) δ 6.81 (s, 1H), 6.66 (s, 1H), 3.87 (s, 3H), 3.76 (d, J= 10 Hz) and 3.29 (t, J=7 Hz) (total 1H, ratio 2:1), 0.95 (t, J=7 Hz, 3H), 0.83 and 0.80 (s, total 3H). FT-IR (neat) 3411, 2923, 1504, 1446, 1267, 1202, 1118, 1024 cm⁻¹. Anal. Calcd for C₂₂H₃₂O₃, 1/4 H₂O: C, 75.71; H, 9.39. Found: C, 75.61; H, 9.33.

EXAMPLE 18

16 β -n-butyl-2-methoxyestradiol

Selected spectral data: ¹H-NMR (300 MHz, CDCl₃) δ 6.81 (s, 1H), 6.66 (s, 1H), 5.43 (s, 1H), 3.88 (s, 3H), 3.76 (d, J=10 Hz) 3.29 (d, J=8 Hz) (total 1H, ratio 2.6:1), 0.83 and 0.80 (s, total 3H). FT-IR (neat) 3221, 2921, 1594, 1504, 1416, 1265, 1200, 1021 cm⁻¹. Anal. Calcd for C₂₃H₃₄O₃: C, 77.04; H, 9.56. Found: C, 77.06; H, 9.65.

EXAMPLE 19

16 β -isobutyl-2-methoxyestradiol

Selected spectral data: ¹H-NMR (300 MHz, CDCl₃) δ 6.81 (s, 1H), 6.66 (s, 1H), 5.43 (s, 1H), 3.88 (s, 3H), 3.77 (dd, J= 9, 10 Hz) and 3.26 (t, J=7 Hz) (total 1 H, ratio 2:1), 0.84 and 0.80 (s, total 3H). IR (neat) 3525, 2913, 1506, 1258, 1202, 1026 cm⁻¹. Anal. Calcd for C₂₂H₃₀O₃: C, 76.69; H, 9.37. Found: C, 76.82; H, 9.47.

EXAMPLE 20

16 β -methyl(dimethyl amine)-2-methoxyestradiol

Selected spectral data: ¹H-NMR (300 MHz, CDCl₃) δ 6.81 (s, 1H), 6.65 (s, 1H), 3.88 (s) and 3.85 (obscured d) (total 4H), 2.28 (s, 6H), 0.87 (s, 3H). Anal. Calcd for C₂₂H₃₃O₃N, 1/4 H₂O: C, 72.59; H, 9.28; N, 3.85. Found: C, 72.80; H, 9.17; N, 3.66.

EXAMPLE 21

16 β -methanol-2-methoxyestradiol

Selected spectral data: ¹H-NMR (300 MHz, CDCl₃) δ 6.78 (s, 1H), 6.61 (s, 1H), 3.92 (d, J=11Hz, 1H), 3.84 (s, 3H), 3.80 (d, J=10 Hz, 1H), 3.63 (d, J=8, 11Hz, 1H), 0.83 (s, 3H). FT-IR (neat)

3283, 3091, 2919, 1602, 1513, 1445, 1204, 1119, 1013 cm^{-1} . Anal. Calcd for $\text{C}_{20}\text{H}_{28}\text{O}_4$: C, 72.25; H, 8.49. Found: C, 72.24; H, 8.48.

EXAMPLE 22

16 α -methanol-2-methoxyestradiol

Selected spectral data: $^1\text{H-NMR}$ (300 MHz, CDCl_3) δ 6.77 (s, 1H), 6.61 (s, 1H), 3.84 (s, 3H), 3.84 (dd, $J=7, 8$ Hz, 1H), 3.61 (dd, $J=9, 11$ Hz, 1H), 3.45 (d, $J=8$ Hz, 1H), 0.83 (s, 3H).

EXAMPLE 23

MDA-MB-231 *In Vitro* Cellular Proliferation Inhibition MDA-MB-231 Cells and Culture Conditions

Figure 1 illustrates the antiproliferative activity in cells and tumor by 2-methoxyestradiol compounds of the present invention which are modified at the 16 position.

MDA-MB-231 human breast carcinoma cells were grown in DMEM containing 10% FCS (Hyclone Laboratories, Logan UT) and supplemented with 2 mM L-Glutamine, 100 units/ml penicillin, 100 $\mu\text{g/ml}$ streptomycin (Irvine Scientific, Santa Anna, CA).

Proliferation Assays

MDA-MB-231 cells were plated at 5000 cells/ml in 96-well plates. After allowing the cells to attach overnight, the appropriate fresh media were applied containing differing concentrations of 2-ME2 or derivatives thereof, as described below. Drug was dissolved in DMSO (Sigma, St. Louis, MO) and added to the wells in a volume of 200 μl . Cells were incubated for two days at 37°C ; at 32 h BrdU was added. BrdU cell proliferation assay (a nucleotide analogue with a fluorescein tag that is incorporated into DNA) was performed as described by the manufacturer (Roche). Each condition was prepared in triplicate and the experiments were carried out a minimum of two times. Results are presented and means \pm SE.

EXAMPLE 24

HUVAC *In Vitro* Cellular Proliferation Inhibition HUVAC Cells and Culture Conditions

HUVAC cells were grown in EGM (Clonetics)

Proliferation Assays

HUVEC cells were plated at 5000 cells/ml in 96-well plates. After allowing the cells to attach overnight, the cells were washed with PBS and incubated in the absence of growth factor for 24 h (EBM, 2% FCS, Clonetics). Cells were treated with increasing concentrations of drug in EBM containing 2% FCS and 10ng/ml bFGF for 48 h at 37°C. Drug preparation, volumes added and BrdU proliferation assay were performed as indicated above.

Results

The breast cancer cell line activities and the cell panels most sensitive to selected analogs are shown in Table 2.

Table 2

R	α / β ratio at position 16	HUVEC IC ₅₀ (μ M)	MDA-MB-231 IC ₅₀ (μ M)
H	N / A	0.5	0.9
methyl (-CH ₃)	All alpha	<0.5	<0.5
methyl	1 : 2	1.3	5
ethyl (-CH ₂ CH ₃)	1 : 1	2	3
n-propyl (-CH ₂ CH ₂ CH ₃)	7.3 : 1	6	>50
n-propyl	1 : 2	9	36
i-butyl $\begin{array}{c} \text{CH}_3 \\ \\ (-\text{CH}_2\text{CHCH}_3) \end{array}$	1 : 2	7.5	40
n-butyl (-CH ₂ CH ₂ CH ₂ CH ₃)	2 : 1	25	82
n-butyl	1 : 2.6	9	39
methanol (-CH ₂ OH)	All alpha	15	22
methanol	All beta	5	50
Methyl-dimethylamine $\begin{array}{c} \text{CH}_3 \\ \\ (-\text{CH}_2\text{NCH}_3) \end{array}$	All beta	9	22

2-Methoxyestradiol is a potent anti-angiogenic and anti-tumor agent. In order to assess the biological activity of modifications at position 16, the anti-proliferative activity of these analogs was evaluated on human umbilical vein endothelial cells (HUVEC) and breast carcinoma cell line, MDA-MB-231 as models for the anti-angiogenic and anti-tumor activity, respectively. It was found that a moderate decrease (approximately 18 fold) in anti-proliferative activity occurs as steric bulk increased (note trend from R=Et to R=Bu). The most active compound in this series is 16 α -methyl, which has greater activity than 2-methoxyestradiol.

The MDA-MB-231 tumor cell line, has a much greater sensitivity to substitutions at position 16 compared to HUVEC cells. Any group at position 16 larger than ethyl has a significant decrease in antiproliferative activity (IC₅₀ >22 μ M). Of the active compounds, 16 α -methyl has better activity than 2-methoxyestradiol,

whereas 16 β -methyl (which is a 1:2 mixture of α : β , so the presence of the α isomer may account for this activity) has about 5-fold less activity than 2-methoxyestradiol, and racemic 16-ethyl has about a 3-fold drop in activity compared to 2-methoxyestradiol.

5 These data suggest that it is possible to design compounds that are selective anti-angiogenic agents. For example, 16 α -propyl is greater than ten-fold less active in inhibiting tumor growth while it has good activity inhibiting endothelial cell proliferation. Other examples include: 16 β -propyl (4-fold difference), 10 16 β -i-butyl (5-fold difference), 16 β -n-butyl (4-fold difference) and 16 β -methanol (10-fold difference). Additionally, a small alkyl group at position 16 can be added without significantly impacting the anti-proliferative activity of the molecule.

15 All of the publications mentioned herein are hereby incorporated by reference in their entireties. The above examples are merely demonstrative of the present invention, and are not intended to limit the scope of the appended claims.
20